Numerical Integration

Computer Graphics
CMU 15-462/15-662

Motivation: The Rendering Equation

- **Recall the rendering equation, which models light "bouncing around the scene":**



$$L_o(\mathbf{p}, \omega_o) = L_e(\mathbf{p}, \omega_o) + \int_{\mathcal{H}^2} f_r(\mathbf{p}, \omega_i \to \omega_o) L_i(\mathbf{p}, \omega_i) \cos\theta \, d\omega_i$$

How can we *possibly* evaluate this integral?

Numerical Integration—Overview

- In graphics, many quantities we're interested in are naturally expressed as integrals (total brightness, total area, …)
- For very, *very* simple integrals, we can compute the solution analytically
- For everything else, we have to compute a numerical approximation
- Basic idea:
  - integral is "area under curve"
  - sample the function at many points
  - integral is approximated as weighted sum

$$\int_0^1 \frac{1}{3}x^2\,dx = \left[x^3\right]_0^1 = 1$$



Rendering: what are we integrating?

- **Recall this view of the world:**

Want to "sum up"—i.e., integrate!—light from all directions

(But let's start a little simpler…)

Review: integral as "area under curve"

$$\int_a^b f(x)dx$$



Or: average value times size of domain

$$\int_a^b f(x)dx = (b-a)\mathrm{mean}(f)$$



Review: fundamental theorem of calculus

$$\int_a^b f(x)dx = F(b) - F(a)$$

$$f(x) = \frac{d}{dx}F(x)$$



Simple case: constant function

$$\int_a^b C dx = (b-a)C$$



Affine function: $f(x) = cx + d$

$$\int_a^b f(x)dx = \frac{1}{2}(f(a) + f(b))(b - a)$$



Need only *one* sample of the function (at just the right place…)

More general polynomials?



Gauss Quadrature

- For any polynomial of degree *n*, we can always obtain the exact integral by sampling at a special set of *n* points and taking a special weighted combination



n=1



n=2



n=4



n=3



n=5

Piecewise affine function

For piecewise functions, just sum integral of each piece:

$$\int_a^b f(x)dx = \frac{1}{2}\sum_{i=0}^{n-1}(x_{i+1} - x_i)(f(x_i) + f(x_{i+1}))$$



Piecewise affine function

If N-1 segments are of equal length: $h = \dfrac{b-a}{n-1}$

$$\int_a^b f(x)dx = \frac{h}{2}\sum_{i=0}^{n-1}(f(x_i) + f(x_{i+1})$$

$$= h\left(\sum_{i=1}^{n-1} f(x_i) + \frac{1}{2}\left(f(x_0) + f(x_n)\right)\right)$$



Key idea so far:

To approximate an integral, we need

(i) quadrature points, and

(ii) weights for each point

$$\int_a^b f(x)\,dx \approx \sum_{i=1}^{n} w_i f(x_i)$$

Arbitrary function f(x)?



Trapezoid rule

<u>Approximate</u> integral of f(x) by assuming function is piecewise linear

For equal length segments: $h = \dfrac{b-a}{n-1}$

$$\int_a^b f(x)dx = h\left(\sum_{i=1}^{n-1} f(x_i) + \frac{1}{2}\left(f(x_0) + f(x_n)\right)\right)$$



Trapezoid rule

Consider cost and accuracy of estimate as $n \to \infty$ **(or** $h \to 0$**)**

Work: $O(n)$

Error can be shown to be: $O(h^2) = O(\frac{1}{n^2})$

(for f(x) with continuous second derivative)



What about a 2D function?



How should we approximate the area underneath?

Integration in 2D

Consider integrating $f(x,y)$ using the trapezoidal rule (apply rule twice: when integrating in *x* and in *y*)

$$\int_{a_y}^{b_y}\int_{a_x}^{b_x} f(x,y)dxdy = \int_{a_y}^{b_y}\left(\boxed{O(h^2)+\sum_{i=0}^{n} A_i f(x_i,y)}\right)dy$$

First application of rule

$$= O(h^2)+\sum_{i=0}^{n} A_i \int_{a_y}^{b_y} f(x_i,y)dy$$

$$= O(h^2)+\sum_{i=0}^{n} A_i \left(\boxed{O(h^2)+\sum_{j=0}^{n} A_j f(x_i,y_j)}\right)$$

Second application

$$= O(h^2)+\sum_{i=0}^{n}\sum_{j=0}^{n} A_i A_j f(x_i,y_j)$$

Errors add, so error still: $O(h^2)$

But work is now: $O(n^2)$

(*n* x *n* set of measurements)

Must perform much more work in 2D to get same error bound on integral!

In K-D, let $N = n^k$

Error goes as: $O\left(\frac{1}{N^{2/k}}\right)$

Curse of Dimensionality

- How much does it cost to apply the trapezoid rule as we go up in dimension?
  - 1D: $O(n)$
  - 2D: $O(n^2)$
  - …
  - kD: $O(n^k)$



- For many problems in graphics (like rendering), *k* is very, very big (e.g., tens or hundreds or thousands)



- Applying trapezoid rule does not scale!
- Need a fundamentally different approach…

Monte Carlo Integration

Credit: many of these slides were created by Matt Pharr and Pat Hanrahan

Monte Carlo Integration

So far we've discussed techniques that use a fixed set of sample points (e.g., uniformly spaced, or obtained by finding roots of polynomial (Gaussian quadrature))

- Estimate value of integral using random sampling of function
  - Value of estimate depends on random samples used
  - But algorithm gives the correct value of integral "on average"

- Only requires function to be evaluated at random points on its domain
  - Applicable to functions with discontinuities, functions that are impossible to integrate directly

- Error of estimate is independent of the dimensionality of the integrand
  - Depends on the number of random samples used: $O(n^{1/2})$

Recall previous trapezoidal rule example: $O(n^{-1/k})$
(dropping the n^2 for simplicity)

Review: random variables

X **random variable. Represents a distribution of potential values**

$X \sim p(x)$ **probability density function (PDF). Describes relative probability of a random process choosing value** x

Uniform PDF: all values over a domain are equally likely

e.g., for an unbiased die

X **takes on values 1,2,3,4,5,6**

$$p(1) = p(2) = p(3) = p(4) = p(5) = p(6)$$

Discrete probability distributions

***n* discrete values** x_i

With probability p_i



Requirements of a PDF:

$$p_i \geq 0$$

$$\sum_{i=1}^{n} p_i = 1$$

Six-sided die example: $p_i = \frac{1}{6}$

Think: p_i **is the probability that a random measurement of** X **will yield the value** x_i

X **takes on the value** x_i **with probability** p_i

Cumulative distribution function (CDF)

(For a discrete probability distribution)



Cumulative PDF: $P_j = \sum_{i=1}^{j} p_i$

where:

$$0 \leq P_i \leq 1$$

$$P_n = 1$$



How do we generate samples of a discrete random variable (with a known PDF?)

Sampling from discrete probability distributions



To randomly select an event, select x_i if

$$P_{i-1} < \xi \leq P_i$$

Uniform random variable $\in [0, 1)$



Continuous probability distributions

PDF $p(x)$

$$p(x) \geq 0$$

CDF $P(x)$

$$P(x) = \int_0^x p(x)\,\mathrm{d}x$$

$$P(x) = \Pr(X < x)$$

$$P(1) = 1$$

$$\Pr(a \leq X \leq b) = \int_a^b p(x)\,\mathrm{d}x$$

$$= P(b) - P(a)$$

Uniform distribution

(for random variable X defined on [0,1] domain)



Sampling continuous random variables using the inversion method

Cumulative probability distribution function

$$P(x) = \Pr(X < x)$$



Construction of samples:

Solve for $x = P^{-1}(\xi)$

Must know the formula for:

1. **The integral of** $p(x)$
2. **The inverse function** $P^{-1}(x)$

Example—Sampling Quadratic Distribution

- As a toy example, consider the simple probability distribution p(x) := 3(1-x)² over the interval [0,1]
- How do we pick random samples distributed according to p(x)?
- First, integrate probability distribution p(x) to get cumulative distribution P(x)
- Invert P(x) by solving y = P(x) for x
- Finally, plug uniformly distributed random values y in [0,1] into this expression



$$\int_0^s 3(1-x)^2\,dx = s^3 - 3s^2 + 3s$$

$$x = 1 - (1-y)^{\frac{1}{3}}$$



How do we uniformly sample the unit circle?



I.e., choose any point P=(px, py) in circle with equal probability)

Uniformly sampling unit circle: first try

- $\theta =$ **uniform random angle between 0 and** 2π
- $r =$ **uniform random radius between 0 and 1**
- **Return point:** $(r \cos\ \theta, r \sin\ \theta)$

This algorithm <u>does not</u> produce the desired uniform sampling of the area of a circle. Why?

Because sampling is not uniform in area!

Points farther from center of circle are less likely to be chosen



$$\theta = 2\pi\xi_1 \qquad r = \xi_2$$

So how should we pick samples? Well, think about how we integrate over a disk in polar coordinates…

Sampling a circle (via inversion in 2D)

$$A = \int_0^{2\pi} \int_0^1 r \, \mathrm{d}r \, \mathrm{d}\theta = \int_0^1 r \, \mathrm{d}r \int_0^{2\pi} \mathrm{d}\theta = \left(\frac{r^2}{2} \right) \Big|_0^1 \theta \Big|_0^{2\pi} = \pi$$

$$p(r, \theta) \, \mathrm{d}r \, \mathrm{d}\theta = \frac{1}{\pi} r \, \mathrm{d}r \, \mathrm{d}\theta \rightarrow p(r, \theta) = \frac{r}{\pi}$$

so that we integrate to 1 instead of area

$$p(r, \theta) = p(r) p(\theta)$$

r, θ independent

$$p(\theta) = \frac{1}{2\pi}$$

$$P(\theta) = \frac{1}{2\pi} \theta \qquad \theta = 2\pi \xi_1$$

$$p(r) = 2r$$

$$P(r) = r^2 \qquad r = \sqrt{\xi_2}$$



Uniform area sampling of a circle

WRONG

probability is uniform; samples are not!



$$\theta = 2\pi\xi_1$$

$$r = \xi_2$$

RIGHT

probability is nonuniform; samples are uniform



$$\theta = 2\pi\xi_1$$

$$r = \sqrt{\xi_2}$$

Uniform sampling via rejection sampling

Completely different idea: pick uniform samples in square (easy)
Then toss out any samples not in square (easy)



Efficiency of technique: area of circle / area of square

Efficiency of Rejection Sampling

- If the region we care about covers only a very small fraction of the region we're sampling, rejection is probably a bad idea:



Smarter in this case to "warp" our random variables to follow the spiral.

Next Time: Monte Carlo Ray Tracing