# Lecture 2: Math (P)Review Part I: Linear Algebra

**Computer Graphics CMU 15-462/15-662** 

### Linear Algebra in Computer Graphics Today's topic: linear algebra.

- Why is linear algebra important for computer graphics?
  - Effective bridge between geometry, physics, etc., and computation.
  - In many areas of graphics, once you can express the solution to a problem in terms of linear algebra, you're essentially done: now ask the computer to solve Ax=b.
  - Fast numerical linear algebra has really made modern computer graphics possible (image processing, physically-based animation, geometry processing...)





## Vector Space—Formal Definition

# Linear algebra is the study of vector spaces and linear maps between them—here's the formal definition\*:

For all vectors **u**, **v**, **w** and scalars *a*, *b*:



### Where do these rules come from?

- In mathematics (and in life) you should never simply accept a set of rules handed to you by an authority...
- Let's try to understand where these "rules" come from.

\*this will NOT be on the test!



### **Vectors - Intuition**

- First things first: what is a vector?
- Intuitively, a vector is a little arrow:



In computer graphics, we work with many types of data that may not look like little arrows (polynomials, images, radiance...). But they still behave like vectors. So, this little arrow is still often a useful mental model.

## **Vectors - What Can We Measure?**

- What information does a vector encode?
- Fundamentally, just direction and magnitude\*:



- For instance, a vector in 2D can be encoded by a length and an angle relative to some fixed direction ("polar coordinates").
- (Side note: are these values the same in any coordinate system?)
- How else might we encode a vector?

\*Traditionally, a vector does not include a "basepoint"; a vector with a basepoint is sometimes called a tangent vector.

### **Vector in Cartesian Coordinates**

# Can also measure components of a vector with respect to some chosen coordinate system:



### René Descartes, Est. 1596

# WARNING: Can't directly compare coordinates in different systems! (Also shouldn't compare (r,θ) to (x,y).)





### What Can We Do with a Vector?

Two basic operations. First, we can add them "end to end":



What if we walk along v first, then u? Actually, it doesn't seem to matter: u+v = v+uLanguage: vector addition is "commutative" or "abelian"



# **Niels Henrik Abel**

### What Else Can We Do with a Vector?

### **Other basic operation? Scaling:**

- In general, can multiply any vector u by a number or "scalar" a to get a new vector au.
- Multiplication behaves the way we would expect, based on the geometric behavior of scaling "little arrows." E.g.,

a(bu) = (ab)u





### **Interaction of Addition & Scaling**

What if we try to add two scaled vectors? Or scale two vectors that have been added together?

 $a(\mathbf{u} + \mathbf{v})$ 





### **Vector Space**—**Formal Definition**

# If we keep playing around vectors, eventually we come up with a complete set of "rules" that vectors seem to obey:

For all vectors **u**, **v**, **w** and scalars *a*, *b*:

• 
$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$$

• 
$$\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$$

- There exists a *zero vector* " $\mathbf{0}$ " such that  $\mathbf{v} + \mathbf{0} = \mathbf{0} + \mathbf{v} = \mathbf{v}$
- For every **v** there is a vector " $-\mathbf{v}$ " such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$
- $1\mathbf{v} = \mathbf{v}$

• 
$$a(b\mathbf{v}) = (ab)\mathbf{v}$$

• 
$$a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$$

• 
$$(a+b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}$$

# *These rules did not "fall out of the sky!"* Each one comes from the geometric behavior of "little arrows." (Can you draw a picture for each one?)

Any collection of objects satisfying all of this properties is a vector space (even if they don't look like little arrows!)

### **Euclidean Vector Space**

- Most common example: Euclidean n-dimensional space
- **Typically denoted by**  $\mathbb{R}^n$ , meaning "n real numbers"
- E.g., (1.23, 4.56,  $\pi/2$ ) is a point in  $\mathbb{R}^3$
- Why such a common example?
  - Looks a lot like the space we live in!
  - That's what we can easily encode on a computer (a list of floating-point numbers).





### **Functions as Vectors**

- Another very important example of vector spaces in computer graphics are spaces of *functions*.
- Why? Because many of the objects we want to work with in graphics are functions! (Images, radiance from a light source, surfaces, modal vibrations, ...)







### Functions as Vectors

Do functions exhibit the same behavior as "little arrows?"
 Well, we can certainly add two functions:

 $\int g(x)$ 



### We can also scale a function:





### **Functions as Vectors**

### What about the rest of these properties?

For all vectors **u**, **v**, **w** and scalars *a*, *b*:

• 
$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$$

- $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- There exists a *zero vector* " $\mathbf{0}$ " such that  $\mathbf{v} + \mathbf{0} = \mathbf{0} + \mathbf{v} = \mathbf{v}$
- For every **v** there is a vector " $-\mathbf{v}$ " such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$

• 
$$1\mathbf{v} = \mathbf{v}$$

• 
$$a(b\mathbf{v}) = (ab)\mathbf{v}$$

• 
$$a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + b\mathbf{v}$$

• 
$$(a+b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}$$

### Try it out at home!

E.g., the "zero vector" is the function equal to zero for all x.

Short answer: yes, functions are vectors! (Even if they don't look like "little arrows".)

## **Vectors in Coordinates**

- So far, we've only drawn our vector operations via pictures.
- How do we actually compute with vectors? **Return to our coordinate representation:**



\*Side note: does it make sense to add vectors encoded as  $(r, \theta)$ ?

# Ok, so we came up with some rule for adding pairs of numbers.

# How can we check that it faithfully encodes geometric behavior of "little arrows?"

### From Geometry to Algebra

### Just check that it agrees with our list of rules that we know (from reasoning *geometrically*) "little arrows" must obey:

For all vectors **u**, **v**, **w** and scalars *a*, *b*:

• 
$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$$

- u + (v + v) (u + v) + w
  There exists a *zero vector* "0" such that v + 0 = 0 + v = v
- For every **v** there is a vector " $-\mathbf{v}$ " such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$
- $1\mathbf{v} = \mathbf{v}$

• 
$$a(b\mathbf{v}) = (ab)\mathbf{v}$$

• 
$$a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$$

• 
$$(a+b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}$$

# For instance, for any two vectors u := (u<sub>1</sub>,u<sub>2</sub>) and v := (v<sub>1</sub>,v<sub>2</sub>) we have

$$\mathbf{u} + \mathbf{v} = (u_1, u_2) + (v_1, v_2) = (u_1 + u_1, v_2) + (v_1, v_2) = (v_1, v_2) + (v_1, v_2$$

### v + 0 = 0 + v = vthat v + (-v) = 0

 $+v_1, u_2 + v_2) =$  $u_1, u_2) = \mathbf{v} + \mathbf{u}.$ 

### **Turning geometric observations into algebraic** rules is convenient for symbolic manipulation & numerical computation.

But you should *never* blindly accept a rule given by authority.

Always ask: where does this rule come from? What does it mean geometrically? (Can you draw a picture?)

### **Scaling Vectors in Coordinates** We'd also like to be able to scale vectors using coordinates. Any ideas?



(From here, check the rest of the properties...)



## **Computing the Midpoint**

- As we start to combine vector operations, we build up operations needed for computer graphics.
- E.g., how would I compute the midpoint m of a = (3,4)and b=(7,2)?



# $m = \frac{1}{2}(a + b)$ $= \frac{1}{2}((3,4) + (7,2))$ $=\frac{1}{2}(10,6)$ = (5,3)

## **Measuring Vectors**

- Earlier we asked, "what information does a vector encode?"
- (A: Orientation and magnitude.)
- How do we actually *measure* these quantities?



## Norm of a Vector

- Let's start with magnitude—for a given vector v, we want to assign it a number |v| called its length or magnitude or norm.
- Intuitively, the norm should capture how "big" the vector is.









CMU 15-462/662

large nor

B

### **Natural Properties of Length—Positivity**

- What properties might you expect the norm (or length) of a vector to satisfy?
- For one thing, it probably shouldn't be negative!

 $|\mathbf{u}| \geq 0$ 

And probably it should be zero only for the zero vector:  $|\mathbf{u}| = 0 \quad \iff \mathbf{u} = \mathbf{0}$ 



## Natural Properties of Length, Continued

Also, if we scale a vector by a factor c, its norm (i.e., length) really should scale by the same amount:



Finally, we know that the shortest path between two points is always along a straight line:



(This final property is sometimes called the "pentagon inequality," since the diagram looks like a pentagon.)



### $|\mathbf{u}| + |\mathbf{v}| \ge |\mathbf{u} + \mathbf{v}|$

### Norm—Formal Definition

A norm is any function that assigns a number to each vector and satisfies the following properties for all vectors u, v, and all scalars a:

• 
$$|\mathbf{v}| \ge 0$$
  
•  $|\mathbf{v}| = 0 \iff$   
•  $|a\mathbf{v}| = |a||\mathbf{v}|$   
•  $|\mathbf{u}| + |\mathbf{v}| \ge |\mathbf{u} + \mathbf{v}|$ 

But you don't have to take my word for it—for each rule, you now have a concrete geometric picture explaining why this "rule" is there.

### $\mathbf{v} = \mathbf{0}$

### **Euclidean Norm in Cartesian Coordinates**

### A standard norm is the so-called *Euclidean norm* of n-vectors:





# natural, geometric properties of a norm?

## L<sup>2</sup> Norm of Functions

- Less familiar idea, but same basic intuition: the so-called L<sup>2</sup> norm measures the total magnitude of a function.
- Consider real-valued functions on the unit interval [0,1] whose square has a well-defined integral. The L<sup>2</sup> norm is defined as:

Not too different from the Euclidean norm: we just replaced a sum with an integral (which is kind of like a sum...).

f(x)



large L<sup>2</sup> norm

small L<sup>2</sup> norm

 $||f|| := \sqrt{\int_0^1 f(x)^2 \, dx}$ 

Q: Careful—does the formula above exactly satisfy all our desired properties for a norm?

### L<sup>2</sup> Norm of Functions—Example

- Consider the function  $f(x) := x\sqrt{3}$ , defined over the unit interval [0,1].
- Q: What is its L<sup>2</sup> norm?

## **A:** $||f||^{2} = \int_{0}^{1} 3x^{2} dx = \left[ x^{3} \right]_{0}^{1} = 1^{3} - 0^{3} = 1$



For clarity we will use  $\|\cdot\|$  for the norm of a

P.S. Most integrals in graphics are not calculated this way (at least not for more challenging functions, or functions described by data). Later on we'll talk a *lot* about <u>numerical integration</u>.



### function, and $|\cdot|$ for the norm of a vector in R<sup>n</sup>.

### **Inner Product**—Motivation

What else can we measure? In addition to magnitude, we said that vectors have *orientation*. Just as norm measured length, inner product measures how well vectors "line up."





(fairly similar)



U



### (quite different!)

### Inner Product—Symmetry

- Will write inner product (also sometimes called the scalar product or dot product) using the notation <u,v> (some folks also write it as u•v).
- When measuring the alignment of two vectors u,v, what are some natural properties you might expect?
- One "obvious" property: order shouldn't matter, since u is just as well-aligned with v as v is with u:



Moreover, simply *re-naming* the vectors should have no effect on how well-aligned they are!

# Inner Product—Projection & Scaling For unit vectors |u|=|v|=1, an inner product measures the extent of one vector along the direction of the other:



If we scale either of the vectors, the inner product also scales:



### **Q:** Is this property symmetric? I.e., is the length of v along u the same as the length of u along v?

 $\langle 2\mathbf{u}, \mathbf{v} \rangle = 2 \langle \mathbf{u}, \mathbf{v} \rangle$ 

### Inner Product—Positivity

Also, a vector should always be aligned with itself, which we can express by saying that the inner product of a vector with itself should be positive (or at least, non-negative): 



In fact, if we continue to think of the inner product of a vector as the length of one vector along another then for unit-length vectors we must have

$$\langle \mathbf{u}, \mathbf{u} \rangle = 1$$

- Q: In general, then, what must <u,u> be equal to? A: Letting  $\hat{u} := u/|u|$ , we have
- $\langle \mathbf{u}, \mathbf{u} \rangle = \langle |\mathbf{u}| \hat{\mathbf{u}}, |\mathbf{u}| \hat{\mathbf{u}} \rangle = |\mathbf{u}|^2 \langle \hat{\mathbf{u}}, \hat{\mathbf{u}} \rangle = |\mathbf{u}|^2 \cdot 1 = |\mathbf{u}|^2$

### $\langle \mathbf{u}, \mathbf{u} \rangle \geq 0$

## Inner Product—Formal Definition

An inner product is any function that assigns to any two vectors u,v a number <u,v> satisfying the following properties:

• 
$$\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$$

• 
$$\langle \mathbf{u}, \mathbf{u} \rangle \geq 0$$

• 
$$\langle \mathbf{u}, \mathbf{u} \rangle = 0 \quad \iff \mathbf{u}$$

• 
$$\langle a\mathbf{u},\mathbf{v}\rangle = a\langle \mathbf{u},\mathbf{v}\rangle$$

•  $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$ 

Q: Which of these properties didn't we talk about? Can you argue that they make sense geometrically? (Discuss online!)





### **Inner Product in Cartesian Coordinates**

# A standard inner product is the so-called *Euclidean* inner product, which operates on a pair of n-vectors:

 $\langle \mathbf{u}, \mathbf{v} \rangle = \langle (u_1, \ldots, u_n), (v_1, \ldots, v_n) \rangle := \sum u_i v_i$ 





u = (4, 1)v = (1, 3)

### $\langle \mathbf{u}, \mathbf{v} \rangle = 4 \cdot 1 + 1 \cdot 3 = 7$

## L<sup>2</sup> Inner Product of Functions—Example

- Just like we had a norm for functions, we can also define an inner product that measures how well two functions "line up".
- E.g., for square-integrable functions on the unit interval:

$$\langle\!\langle f,g\rangle\!\rangle := \int_0^1 f(x)g$$



(x) dx

# Measuring Images, Other Signals?

- Many ways to measure "how big" a signal is (norm) or "how well-aligned" two signals are (inner product).
- Choice of inner product depends on application.
- For instance, suppose we want images with "interesting stuff"
- Might try measuring norm of *derivative* (captures edges):









## Linear Maps

- At the beginning, said linear algebra was study of vector spaces and linear maps between them.
- Have a pretty good handle on vector (and inner product) spaces.
- But what's a linear map? And why is it useful for graphics?
- We'll get to the 1st question in a moment. As for the 2nd question, a few reasons:
  - Computationally, easy to solve systems of linear equations.
  - Basic transformations (rotation, translation, scaling) can be expressed as linear maps. (Will see this in a later lecture!)
  - All maps can be approximated as linear maps over a short distance/short time. (Taylor's theorem). This approximation is used all over geometry, animation, rendering, image processing...

## Linear Maps—Geometric Definition

- What is a linear map?
- Especially in graphics, can think about them visually.
- **Example:**



### **Key idea:** *linear maps take lines to lines*\*



### Linear Maps—Geometric Definition

- What is a linear map?



### **Key idea:** *linear maps take lines to lines*\*

\*...while keeping the origin fixed.



### Linear Maps—Algebraic Definition

A map f is linear if it maps vectors to vectors, and if for all vectors u,v and scalars a we have

$$f(\mathbf{u} + \mathbf{v}) = f(\mathbf{u}) + f(\mathbf{u}) = af(\mathbf{u})$$

In other words: if it doesn't matter whether we add the vectors and then apply the map, or apply the map and then add the vectors (and likewise for scaling):





### **Linear Maps in Coordinates**

- For maps between R<sup>m</sup> and R<sup>n</sup> (e.g., a map from 2D to 3D), we can give an even more explicit definition.
- A map is linear if it can be expressed as

$$f(u_1,\ldots,u_m)=\sum_{i=1}^m u_i\mathbf{a}_i$$

In other words, if it is a linear combination of a fixed set of vectors a<sub>i</sub>:



### Q: Is f(x) := ax + b a linear function?



However, it's not a line through the *origin*, i.e.,  $f(0) \neq 0$ . Another way to see it's not linear? Doesn't preserve sums:

$$f(x_1 + x_2) = a(x_1 + x_2) + b = a$$
$$f(x_1) + f(x_2) = (ax_1 + b) + (ax_2 + b) = a$$

- This function is called an AFFINE function (not a LINEAR one).
- Later we'll see an important computer graphics magic trick: turn affine functions (e.g., translation) into linear ones via homogeneous coordinates.

# $ax_1 + ax_2 + b$ $= ax_1 + ax_2 + 2b$

### More interesting question:

Q: Is  $f(u) := \int_0^1 u(x) dx$  a linear map?

# (Think about it — it will be part of your homework!)

## Span

Q: Geometrically, what is the span of two vectors u, v?

A: The span is the set of all vectors that can be written as a linear combination of u and v, i.e., vectors of the form



### **Span & Linear Maps**

- Just a bit of language—can connect "span" and "linear map":
- "The image of any linear map is the span of some collection of vectors."



Q: What's the image of a function?

### Basis

- Span is also closely related to the idea of a *basis*.
- In particular, if we have exactly n vectors e<sub>1</sub>, ..., e<sub>n</sub> such that

$$\operatorname{span}(\mathbf{e}_1,\ldots,\mathbf{e}_n) =$$

- Then we say that these vectors are a basis for R<sup>n</sup>.
- Note: many different choices of basis!
- Q: Which of the following are bases for the 2D plane (n=2)?



 $= \mathbb{R}^n$ 

### **Orthonormal Basis**

- Most often, it is convenient to have to basis vectors that are (i) unit length and (ii) mutually orthogonal.
- In other words, if  $e_1, \ldots, e_n$  are our basis vectors then

$$\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \begin{cases} 1, & i = j \\ 0, & \text{otherwise} \end{cases}$$

This way, the geometric meaning of the sum  $u_1^2 + \ldots + u_n^2$  is maintained: it is the length of the vector u.



se.

**Common bug: projecting** a vector onto a basis that is **NOT orthonormal while** continuing to use standard norm / inner product.

### **Gram-Schmidt**

- Given a collection of basis vectors  $a_1, \ldots a_n$ , how do we find an orthonormal basis  $e_1, \ldots e_n$ ?
- **Gram-Schmidt algorithm:** 
  - normalize the first vector (i.e., divide by its length)
  - subtract any component of the 1st vector from the 2nd one
  - normalize the 2nd one
  - repeat, removing components of first k vectors from vector k+1



\*WARNING: for large number of vectors / nearly parallel vectors, not the best algorithm...

## **Fourier Transform**

- Functions are also vectors. Do they have an orthonormal basis?
- Yes! This is the basic idea behind the *Fourier transform*.
- Simple example: functions that repeat at intervals of  $2\pi$ :



**Can project onto basis of sinusoids:**  $\cos(nx)$ ,  $\sin(mx)$ ,  $m, n \in \mathbb{N}$ - really just a linear map from one basis to another – fundamental building block for many graphics algorithms



lots of low- and mid-frequency oscillation

not as much high-frequency oscillation

## **Frequency Decomposition of Signals**

- More generally, this idea of projecting a signal onto different "frequencies" is known as Fourier decomposition
- Can be applied to all sorts of signals; basic tool used across, image processing, rendering, geometry, physical simulation...
- Will have plenty more to say as course goes on!





## **System of Linear Equations**

- A system of linear equations is exactly what it sounds like: a bunch of equations where left-hand side is a linear function, right hand side is constant. E.g.,
  - x + 2y = 34x + 5y = 6
- Unknown values are sometimes called "degrees of freedom" (DOFs); equations sometimes called "constraints"
- Goal: solve for DOFs that simultaneously satisfy constraints:

$$x = 3 - 2y$$
$$4(3 - 2y) + 5y = 6$$
$$\boxed{y = 2}$$

### What does solving a linear system mean?

### Linear System, Visualized

- Of course, a linear system can be used to represent many different practical tasks (simulation, processing, etc.).
- But for any linear system, there are some good mental models to visualize:



# e good mental models

# GIVEN a point b, FIND the point x that maps to it:

 $f(\mathbf{x})$ 

### Uniqueness, Existence of Solutions

Of course, not all linear systems can be solved! (And even those that can be solved may not have a unique solution.) 



ax + by = c

### MANY SOLUTIONS any point along the line

### Wait, what about matrices?!

# Matrices in Linear Algebra

- Linear algebra often taught from the perspective of *matrices*, i.e., pushing around little blocks of numbers.
- But linear algebra is not fundamentally *about* matrices.
- As you've just seen, you can understand almost all the basic concepts without ever touching a matrix!
- Likewise, matrices can interfere with understanding / lead to confusion, since the same object (a block of numbers) is used to represent many different things (linear map, quadratic form, ...) in many different bases.
  - Still, VERY useful!
    - symbolic manipulation
    - numerical computation



### **Representing Linear Maps via Matrices** Key example: suppose I have a linear map $\mathbb{R}^2$

$$f(\mathbf{u}) = u_1 \mathbf{a}_1 + u_2 \mathbf{a}_2$$



- How do I encode as a matrix?
- Easy: "a" vectors become matrix columns:

$$A := \begin{bmatrix} a_{1,x} & a_{2,x} \\ a_{1,y} & a_{2,y} \\ a_{1,z} & a_{2,z} \end{bmatrix}$$

Now, matrix-vector multiply recovers original map:

$$\begin{bmatrix} a_{1,x} & a_{2,x} \\ a_{1,y} & a_{2,y} \\ a_{1,z} & a_{2,z} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} a_{1,x}u_1 + a_{2,z} \\ a_{1,y}u_1 + a_{2,z} \\ a_{1,z}u_1 + a_{2,z} \end{bmatrix}$$

 $_{\chi}u_{2}$  $y u_2$  $= u_1 \mathbf{a}_1 + u_2 \mathbf{a}_2$  $_{\chi}\mathcal{U}_{2}$ 

# Don't worry: if you love matrices, there will be plenty of them in your homework!

### P.S. What's the "Pentagon inequality?"



### **Clearly not a pentagon... ASK QUESTIONS!**

## Next time: Math (P)Review Part II

### Vector calculus

- Eigenvalue problems
- Complex numbers

 $\varphi \qquad X$ 

grad  $\phi$ 

div X



### curl Y